

Differences in carbon and nitrogen abundances between field and cluster early-type galaxies

Sánchez-Blázquez P., Gorgas J., Cardiel N.¹ and Cenarro J.

*Departamento de Astrofísica, Facultad de Físicas, Universidad Complutense de Madrid,
28040 Madrid, Spain*

pat@astrax.fis.ucm.es

González J.J.

*Inst. de Astronomía, Universidad Nacional Autónoma de México, Apdo-Postal 70-264,
México D.F, México*

ABSTRACT

Central line-strength indices were measured in the blue spectral region for a sample of 98 early-type galaxies in different environments. For most indices (Mgb and $\langle\text{Fe}\rangle$ in particular) ellipticals in rich clusters and in low-density regions follow the same index-sigma relations. However, striking spectral differences between field ellipticals and their counterparts in the central region of the Coma cluster are found for the first time, with galaxies in the denser environment showing significantly lower C4668 and CN₂ absorption strengths. The most convincing interpretation of these results is a difference in abundance ratios, arising from a distinct star formation and chemical enrichment histories of galaxies in different environments. An scenario in which elliptical galaxies in clusters are fully assembled at earlier stages than their low-density counterparts is discussed.

Subject headings: galaxies: abundances — galaxies: evolution — galaxies: clusters — galaxies: formation — galaxies: stellar content

1. Introduction

Despite the observational and theoretical efforts of the last decades, the evolutionary status of early-type galaxies is still an unsolved problem. The stellar populations of nearby

¹Also at Calar Alto Observatory, CAHA, Apdo. 511, 04044, Almería, Spain

ellipticals preserve a record of their formation and evolution. In particular, the study of their element abundance ratios should be a powerful discriminant between different star formation histories (e.g. Worthey 1998). However, this last approach is still at its infancy. The pioneering works of late 1970s already revealed that abundance ratios in early-type galaxies are often non-solar (O’Connell 1976; Peterson 1976). Since then, several studies have provided compelling evidence of Mg/Fe overabundances in massive ellipticals as compared with the solar ratio (Worthey, Faber, & González 1992; Peletier 1989; Vazdekis et al. 1997), which have been interpreted in the light of several possible scenarios based on the understanding that Mg is mainly produced in type II supernovae (Faber, Worthey, & González 1992; Worthey, Faber, & González 1992; Matteucci 1994). However, and in contrast with the above findings, another *alpha* element, namely Ca, seems to be underabundant in ellipticals (O’Connell 1976; Saglia et al. 2002; Cenarro et al. 2003; Thomas, Maraston, & Bender 2003), challenging current chemical evolution models (Matteucci 1994; Mollá & García-Vargas 2000).

Several authors (Worthey 1998; Vazdekis et al. 2001) have also noted a strengthening in other absorption line-strengths, in particular in the IDS/Lick C4668 and CN₂ indices, when compared with stellar population models predictions. The variations of these two indices are mainly driven by C and N (in the case of CN) abundances (Tripicco & Bell 1995), which could suggest a possible enhancement of these two elements relative to Fe when compared with the solar values. In contrast with Mg, C and N are mainly produced in low- and intermediate-mass stars (Renzini & Voli 1981; Chiappini, Romano, & Matteucci 2003, although there are recent suggestions that most of the C should come from massive stars). During the AGB phase, these stars eject into the ISM significant amounts of ⁴He, ¹²C, ¹³C and ¹⁴N, enriching the medium from which new stars will be formed. Therefore, it seems difficult to simultaneously reproduce the abundances of all these elements with a simple chemical evolution scenario.

The problem of C and N abundances has been more thoroughly studied in the field of globular clusters. An interesting puzzling problem is the existence of a CN dichotomy in Galactic and M31 globular clusters (Burstein et al. 1984). Although this is a controversial issue, recent works (Harbeck, Smith, & Grebel 2003) tend to favor the scenarios of different abundances in the parental clouds against the ones that predict abundance changes produced internally by the evolution of the stars (see Cannon et al. 1998 for a review).

Given the expected sensitivities of relative abundances to the star formation history of ellipticals, their study in galaxies within different environments should help to discriminate between different formation and evolution models. For instance, hierarchical scenarios predict that ellipticals in rich clusters assembled completely at high redshift ($z > 3$), whereas field ellipticals may have experienced an elapsed and more complex star formation history

(Kauffmann & Charlot 1998). However, very little is known about the dependence of the relative abundances on environment. One piece of information is that there is no difference in the $[\text{Mg}/\text{Fe}]$ ratio between cluster and field elliptical galaxies (Jørgensen 1999; Kuntschner et al. 2002). In this letter, we study the behaviour of several Lick/IDS indices (see Worthey et al. 1994 for definition) in a sample of low and high density environment galaxies (LDEG and HDEG respectively) and, surprisingly, we do find systematic differences in the strength of C and CN features.

2. Observations and data analysis

2.1. Observations

Long-slit spectra of 98 early-type galaxies in different environments were taken in four observing runs with two different telescopes. The sample comprises 59 galaxies from the field and the Virgo cluster (LDEG), and 34 galaxies from the central region of the Coma cluster (HDEG), spanning a wide range of absolute magnitudes ($-22.5 < M_B < -16.5$, using $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$), and central velocity dispersions $40 < \sigma < 400 \text{ km s}^{-1}$ (from dwarf ellipticals, in Virgo and Coma, to giant galaxies).

In the first two runs (1998 January and 1999 August) we used the 3.5m telescope at Calar Alto Observatory (Almería, Spain), employing the Twin Spectrograph. The observations of the third and fourth runs (1999 March and 2001 April) were carried out with the 4.2m WHT at the Roque de los Muchachos Observatory (La Palma, Spain) using the ISIS spectrograph. Spectral resolutions range from $2.6 \text{ \AA}$ and $4.0 \text{ \AA}$ (FWHM) for LDEG to $8.6 \text{ \AA}$ for HDEG, in a spectral range $\lambda\lambda 3600\text{--}5400 \text{ \AA}$. Exposures times of 1200–3600 secs per galaxy allowed us to obtain central spectra with signal-to-noise (S/N) ratios (per $\text{\AA}$) ranging from 25 to 250. We also observed several galaxies in common between runs to ensure that the measurements were in the same system. 85 stars from the IDS/Lick library were included to transform the measured line-strength indices to the Lick system.

Standard data reduction procedures (flat-fielding, cosmic ray removal, wavelength calibration, sky subtraction and fluxing) were performed with REDUCE (Cardiel 1999), which allowed a parallel treatment of data and error frames and provided an associated error file for each individual data spectrum.

2.2. Velocity dispersion and line-strength indices

For each galaxy, central spectra were extracted by coadding within a standard metric aperture size corresponding to 4 arcsec projected at the distance of the Coma cluster (simulating in this way a fixed linear aperture of length 1.8 kpc in all the galaxies). Velocity dispersions were determined using the MOVE and OPTIMA algorithms described in González (1993). We measured the Lick/IDS line-strengths indices, although only CN_2 , Mgb, C4668 and $\langle \text{Fe} \rangle$ are presented here (the rest of the indices, with a more detailed explanation of data handling, will be presented in a forthcoming paper). All the indices were transformed to the Lick spectrophotometric system using the observed stars and following the prescriptions in J. Gorgas, P. Jablonka, & P. Goudfrooij (2003, in preparation); see also Worthey & Ottaviani (1997). Using galaxies in common with Trager (1997) and with repeated observations between runs, we double-checked that there are no systematic errors in the indices of galaxies observed in different runs. All the indices presented in this work are transformed into magnitudes, following Kuntschner (1989): $I'(\text{mag}) = -2.5 \log(1 - I(\text{\AA})/\Delta\lambda)$, where $\Delta\lambda$ is the width of the index bandpass.

2.3. Results

In Fig. 1 we present the CN_2 , C4668', Mgb', and $\langle \text{Fe} \rangle'$ indices versus velocity dispersion (σ) for the 98 galaxies of the sample. The spectra of three dwarf galaxies from the Coma sample did not have enough S/N to derive a reliable measurement of σ and a typical value of 40 km s^{-1} was assumed. Also, for the rest of the galaxies with $\sigma < 60 \text{ km s}^{-1}$, we adopted σ values from Pedraz et al. (2002) and Guzmán et al. (2003). It is apparent from this figure that galaxies located in low and high density environments show systematic differences in C4668' and CN_2 , being the indices in HDEG systematically lower than in LDEGs. On the other hand, both galaxy subsamples seem to follow similar relationships in the $\langle \text{Fe} \rangle'$ and Mgb' versus σ diagrams.

To quantify the possible systematic differences, we have performed a linear least-square fit to the LDEG subsample, and have measured the mean offsets (weighting with errors) of the Coma galaxies (HDEG) from the fits. These differences and their errors are indicated within each panel, confirming the high significance of the systematic offsets in CN_2 and C4668'.

These systematic differences are also visible directly in the spectra of the galaxies. For illustration, Fig. 2 shows the coadded spectra of LDEG (thin line) and HDEG (thick line) in the range $150 < \sigma < 250 \text{ km s}^{-1}$, previously shifted to the same radial velocity and

broadened to the maximum σ . The S/N ratios of the two combined spectra are above 300. It is evident that the offsets found in Fig. 1 are due to real enhancements of the CN band at $\lambda 4177 \text{ \AA}$ and the C_2 Swan band at $4735 \text{ \AA}$ in LDEG compared to HDEG. Note that this effect is also quite remarkable in the CN band around $\lambda 3865 \text{ \AA}$ (not included in the Lick system).

3. Discussion

Prior to interpret the systematic differences in CN_2 and $C4668'$ as variations on elements abundances, and since the indices are also sensitive to other physical parameters, we have explored further possibilities:

(1) Given the gravity dependences of the CN_2 and $C4668'$ indices (both are stronger in giant than in dwarf stars; Worthey et al. 1994), a decrease in the giant/dwarf ratio in HDEG (with respect to LDEG) would lead to lower index values. Using the models by Vazdekis et al. (1996), we have checked that a change in the exponent of the IMF from 0.80 to 2.80 would decrease CN_2 and $C4668'$ by 0.033 and 0.009 mag respectively, while the expected changes in Mgb' and $\langle Fe \rangle$ would be of 0.005 and 0.003 in the opposite sense (assuming an age of 10 Gyr and solar metallicity). These predictions are marginally consistent with our results (note that the above offsets for Mgb' and $\langle Fe \rangle$ are compatible with the measurements within the uncertainties), thus we cannot reject this possibility. However, this result is in contradiction with other studies (Rose et al. 1994) which suggest a decrease in the giant/dwarf ratio in LDEG compared to HDEG, and would imply important changes in other observables. Measurements of the Ca triplet in the near-infrared (with a high sensitivity to the IMF; see Cenarro et al. 2003) should help to discard or confirm this scenario.

(2) A difference in (luminosity-weighted) mean age between HDEG and LDEG could also introduce systematic offsets in CN_2 and $C4668'$ between both galaxy samples. Under this scenario, and using the models by V96, the observed offsets could be accounted for if HDEG were about 8 Gyr younger than LDEG. This possibility can be rejected since it would imply a decrease in Mgb' and $\langle Fe \rangle$ of 0.026 and 0.014 in HDEG compared with LDEG, which is not observed. Furthermore, it contradicts previous findings which suggest that HDEG are, in any case, older than LDEG (Kuntschner et al. 2002).

Thus, the most plausible explanation of our results is that variations in the relative abundances of C and N with respect to Mg and Fe are responsible for the observed offsets between galaxies in different environments. Fig. 3 compares the observed CN_2 and $C4668'$ with the predictions of stellar population models. This figure clearly shows that the previ-

ously found overabundances of C and N only stand for LDEG, while HDEG tend to exhibit relative abundances closer to the solar partition.

The C4668' index is extremely sensitive to carbon changes, so little variations in carbon abundance can change this index dramatically (Tripicco & Bell 1995). However, the variations of CN are mostly controlled by N, because extra C is readily incorporated into CO but extra N makes more CN molecules. Therefore, a change in both C and N abundances is required to explain our results. Besides, if extra carbon is easily incorporated into CO, one should detect an enhancement in the strength of the CO bands in LDEG compared to HDEG. This effect has indeed been found by Mobasher & James (1996) comparing the CO band at $2.3\mu\text{m}$ in galaxies from the field and from the Pisces and Abell 2634 clusters. They interpreted this difference as an evidence of intermediate-age stellar population in LDEG (through a major contribution of AGB stars). Although we do not discard a larger contribution of younger populations in LDEG compared to HDEG, our results in the blue spectral range imply that their observations can be solely explained by a relative abundance difference. In any case, the observed offsets can not be due to an age effect alone (see above).

The most plausible scenario to explain the differences in relative abundances is the one in which LDEG and HDEG have experienced different star formation histories. In particular, since the ISM is progressively enriched in C and N by stars of low and intermediate mass stars, HDEG should have been fully assembled before the massive release of these elements. The hierarchical clustering paradigm currently predicts that galaxies in clusters formed at different epochs than LDEGs. If the time elapsed between the assembling of the former and the later is enough to permit the C and N enrichment of the ISM of the pre-merging building blocks in LDEG, the stars formed in these galaxies during the merging events will be C and N enhanced. The constancy of the iron-peak elements could be understood if, in both environments, the mergers take place before type-I SN can significantly pollute the ISM of the pre-merging blocks. Additionally, HDEG could have experienced a truncated star formation and chemical enrichment history compared to a more continuous time-extended history for their counterparts in low density environments. However, under this hypothesis, there should be an increase of magnesium (produced by type-II SN) in LDEG which is not detected. One way to understand the constancy of the Mgb' index could be to invoke the IMF-metallicity relationship suggested by Cenarro et al. (2003), in which the successive episodes of star formation in LDEG would take place with lower giant-to-dwarf ratios.

To conclude, we have noted for the first time a systematic difference in the element abundance ratios of galaxies situated in different environments. These differences impose strong constraints to models of chemical evolution and galaxy formation. It is clear that more work is still needed to completely understand the causes of the differences. In particular, it

would be very interesting to compare the CN_2 and C4668' indices between the central regions and the outskirts of the Coma cluster, where Mobasher & James (2000) found differences in the strength of the near-IR CO molecule. Also, the detailed study of other dense clusters is highly needed to confirm whether this effect is particular of the Coma cluster.

Acknowledgments

We are indebted to R. Guzmán for providing us with a catalogue of dwarf elliptical galaxies in the Coma cluster. The WHT is operated on the island of La Palma by the Royal Greenwich Observatory at the Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias. The Calar Alto Observatory is operated jointly by the Max-Planck-Institute für Astronomie, Heidelberg, and the Spanish Comisión Nacional de Astronomía. This work was supported by the Spanish research project No.AYA2000-977.

REFERENCES

- Burstein, D., Faber, S., Gaskell, C. M., & Krumm, N. 1984, *ApJ*, 287, 586
- Cannon, R. D., Croke, B. F. W., Bell, R. A., Hesser, J. E, & Stathakis, R. A. 1998, *MNRAS*, 298, 601
- Cardiel, N. 1999, PhD thesis, Universidad Complutense, Madrid
- Cenarro, A. J., Gorgas, J., Vazdekis, A., Cardiel, N., & Peletier, R. F. 2003, *MNRAS*, 339, L12
- Chiappini, C., Romano, D., & Matteucci, F. 2003, *MNRAS*, 339, 63
- Faber, S. M., Worthey, G., & González, J. J. 1992, in *IAU Symp. 149, Stellar Populations of Galaxies*, ed. B . Barbuy ,& A. Renzini (Dordrecht: Kluwer), 255
- González, J. J. 1993, PhD thesis, University of California, Santa Cruz.
- Guzmán, R., Graham, A., Matkovic, A., Vass, I., Gorgas, J., & Cardiel, N. 2003, preprint (astro-ph/0303390)
- Harbeck, D., Smith, G., & Grebel, E. 2003, *AJ*, 125, 197
- Jørgensen I. 1999, *MNRAS*, 306, 607
- Kauffmann, G., & Charlot, S. 1998, *MNRAS*, 294, 705
- Kuntschner, H. 1989, PhD thesis, University of Durham, UK.
- Kuntschner, H., Smith, R., Colles, M., Davies, R., Kaldare, R., & Vazdekis, A. 2002, *MNRAS*, 337,172
- Matteucci, F. 1994, *A&A*, 288, 57
- Mobasher, B., & James, P. A. 1996, *MNRAS*, 280, 895
- Mobasher, B., & James, P. A. 2000, *MNRAS*, 316, 507
- Mollá, M., & García-Vargas, M. L. 2000, *A&A*, 359, 18
- O’Connell, R. W. 1976, *ApJ*, 206, 370
- Pedraz, S., Gorgas, J., Cardiel, N., Sánchez-Blázquez, P., & Guzmán, R. 2002, *MNRAS*, 332, 59

- Peletier, R. F. 1989 PhD thesis, Univ. of Groningen
- Peterson, R. C. 1976, *ApJ*, 210, L123
- Renzini, A., & Voli, M. 1981, *A&A*, 94, 175
- Rose, J. A., Bower, R., Caldwell, N., Ellis, R., Sharples, R. M., & Teague, P. 1994, *AJ*, 108, 205
- Saglia, R. P., Maraston, C., Thomas, D., Bender, R., & Colles, M. 2002, *ApJ*, 579, L13
- Thomas, D., Maraston, C., & Bender, R. 2003, *MNRAS*, in press (astro-ph/0303615)
- Trager, S. C. 1997, PhD thesis, University of California, Santa Cruz.
- Tripicco, M. J., & Bell, R. A. 1995, *AJ*, 110, 3035
- Vazdekis, A., Casuso, E., Peletier, R. F., & Beckman, E. 1996, *ApJS*, 106, 307
- Vazdekis, A., Peletier, R. F., Beckman, J.E & Casuso, E. 1997, *ApJS*, 111, 203
- Vazdekis, A., Kuntschner, H., Davies, R., Arimoto, N., Nakamura, O., & Peletier, R. 2001, *ApJ*, 551, 127
- Worthey, G. 1998, *PASP*, 110, 888
- Worthey, G., & Ottaviani, D. L. 1997, *ApJS*, 111, 377.
- Worthey, G., Faber, S. M., & González, J. J. 1992, *ApJ*, 398, 69
- Worthey, G., Faber, S. M., González, J. J., & Burstein, D. 1994, *ApJS*, 94, 687




Fig. 1.— Relations of the line-strength indices analyzed in this paper with central velocity dispersion for HDEG and LDEG. LDEG are represented with open symbols whereas filled symbols correspond to galaxies from the Coma cluster. Dwarfs ellipticals are plotted with triangles, S0 with squares and E galaxies with circles. Lines represent error-weighted least-squares linear fits to the LDEG subsample. Typical errors in the indices and σ 's are included at the bottom-left of each panel. Labels indicate the mean offsets, and their corresponding errors, of the HDEG with respect to the fits.




Fig. 2.— Coadded spectra for LDEG and HDEG, represented with thin and thick lines respectively, in our spectral range. See text for details. The position of the CN band around $\lambda 4177$ Å, and the central bandpasses of the CN₂ and C4668' indices are indicated. Besides these bands, a very consistent difference in the CN violet system (3850–3890 Å) is also evident.




Fig. 3.— C4668' line-strengths versus CN_2 for the complete samples of HDEG and LDEG. Overplotted are models by Vazdekis (1996). Lines of constant $[\text{Fe}/\text{H}] = -1.68, -0.68, -0.38, 0.0, +0.2$ are shown by solid lines. Dashed lines represent models of constant ages, from 1 Gyr to 17.78 Gyr, increasing from left to right. Symbols are the same as in Fig. 1.